

Table 5a. Reported Total Early Syphilis (TES)* Cases^ by Race and Select Demographics, Year to Date^

| | Asian/Pacific Islander | | Black | | Hispanic | | White | | Other/Not Reported | | Total | |
|--------------------------------|---------------------------|-------------|------------|-------------|-----------|-------------|------------|-------------|--------------------|-------------|------------|-------------|
| <u>Sex</u> | | | | | | | | | | | | |
| Female | 0 | 0% | 33 | 13% | 0 | 0% | 9 | 8% | 0 | 0% | 42 | 11% |
| Male | 7 | 100% | 216 | 86% | 16 | 100% | 98 | 91% | 2 | 100% | 339 | 88% |
| All | 7 | 100% | 249 | 100% | 16 | 100% | 107 | 100% | 2 | 100% | 381 | 100% |
| <u>Age at diagnosis</u> | | | | | | | | | | | | |
| 15 - 19 | 0 | 0% | 27 | 10% | 1 | 6% | 1 | 0% | 0 | 0% | 29 | 7% |
| 20 - 29 | 1 | 14% | 117 | 46% | 4 | 25% | 26 | 24% | 1 | 50% | 149 | 39% |
| 30 - 39 | 4 | 57% | 52 | 20% | 6 | 37% | 20 | 18% | 0 | 0% | 82 | 21% |
| 40 - 49 | 1 | 14% | 40 | 16% | 3 | 18% | 36 | 33% | 1 | 50% | 81 | 21% |
| 50+ | 1 | 14% | 13 | 5% | 2 | 12% | 24 | 22% | 0 | 0% | 40 | 10% |
| All | 7 | 100% | 249 | 100% | 16 | 100% | 107 | 100% | 2 | 100% | 381 | 100% |
| <u>Region</u> | | | | | | | | | | | | |
| CENTRAL | 1 | 14% | 93 | 37% | 1 | 6% | 37 | 34% | 2 | 100% | 134 | 35% |
| EASTERN | 1 | 14% | 102 | 40% | 2 | 12% | 22 | 20% | 0 | 0% | 127 | 33% |
| NORTHERN | 5 | 71% | 33 | 13% | 13 | 81% | 33 | 30% | 0 | 0% | 84 | 22% |
| NORTHWEST | 0 | 0% | 3 | 1% | 0 | 0% | 10 | 9% | 0 | 0% | 13 | 3% |
| SOUTHWEST | 0 | 0% | 18 | 7% | 0 | 0% | 5 | 4% | 0 | 0% | 23 | 6% |
| All | 7 | 100% | 249 | 100% | 16 | 100% | 107 | 100% | 2 | 100% | 381 | 100% |

* Total Early Syphilis includes: Primary, Secondary and Early Latent Syphilis

^ See definitions under [Technical Notes](#)

Reported Cases of TES by Region, Year to Date

